

Guidelines for the Care of People with Spina Bifida

Physical Activity

Workgroup Members: Kerri A Vanderbom, PhD, (Chair); John Foley, PhD; David Kanter, MD; Lynne Romeiser Logan, PT, PhD, PCS; Amy C McPherson, PhD

We thank the work of the 2018 workgroup members: Kerri A Vanderbom, PhD; John Foley, PhD; David Kanter, MD; Lynne Romeiser Logan, PT, PhD, PCS; Amy C McPherson, PhD

Introduction

The benefits of physical activity are well documented in the 2018 Physical Activity Guidelines Advisory Committee Scientific Report by the U.S. Department of Health and Human Services.¹² Regular engagement in physical activity has been associated with lower rates of heart disease, depression, type II diabetes, and stroke². It has been suggested that time spent in physical activity should be used as a vital sign³ because it is a leading indicator of health and lowers mortality and morbidity.⁴ Unfortunately, the majority of Americans fail to meet the recommended guidelines for physical activity.⁵ Additionally, it is recognized that both children and adults with Spina Bifida, as well as groups of individuals with other disabilities, are even less active when compared to the general population.^{6–8}

Increasing physical activity for individuals with Spina Bifida may be critical since a loss of strength or fitness may lead to less independence and function in carrying out activities of daily living.⁹ In addition to physical and functional benefits of physical activity, there are many mental health benefits such as perceived improvement in activities of daily living¹⁰ higher athletic competence, better perceived physical appearance, greater self-worth, and higher quality of life.¹¹

The World Health Organization (WHO) has updated its global recommendations on physical activity and sedentary behavior, reinforcing that any movement is better than none and that reducing sedentary time is as important as increasing physical activity.^{12,13} The updated guidelines emphasize the importance of tailoring physical activity recommendations to individuals with disabilities, including those with Spina Bifida, to promote overall health and prevent secondary complications¹⁴. According to the WHO, individuals with disabilities should engage in regular physical activity within their abilities, incorporating aerobic, muscle-strengthening, and flexibility exercises to maintain function and prevent deconditioning.¹² It should be noted that there is still debate on how much physical activity is enough for persons with certain disabilities, including individuals with impairments.^{15,16} There is agreement on the importance of reducing sedentary behavior and aiming to replace it with activity, but the question is how much physical activity? some argue that there is not enough evidence to strongly suggest that the physical activity recommendations for persons without disabilities have the same benefit for all individuals.^{15,17}

That said, research highlights that individuals with Spina Bifida can achieve improved health outcomes through structured physical activity interventions. Studies have shown that aerobic and resistance training can significantly enhance cardiorespiratory endurance, muscle strength, and functional mobility in individuals with Spina Bifida.¹⁸ Participation in leisure sports is also associated with higher bone mineral density and reduced obesity risk.¹⁹ Furthermore, technology-assisted

Guidelines for the Care of People with Spina Bifida

interventions, such as virtual reality (VR) and exergaming, have demonstrated effectiveness in engaging individuals with Spina Bifida in physical activity while improving motivation and adherence.²⁰ These findings suggest that a combination of structured, community-based, and technology-driven approaches may help reduce the high levels of sedentary behavior commonly observed.^{21,22}

The National Physical Activity Guidelines are for everybody.²³ These guidelines are incorporated into the Spina Bifida Guidelines for Physical Activity and should be followed as closely as possible by children and adults with Spina Bifida unless deemed medically unsafe by a health care provider. Physical activity is the key to health and wellness. Endorsement rather than caution is recommended for the vast majority of individuals with Spina Bifida.

The physical activity guidelines for children ages 6-17 state:²³

- Children should engage in 60 minutes or more of physical activity each day.
- Aerobic activity should make up most of the youth's activity each day; vigorous intensity aerobic activity should be done at least 3 days/week.
- Muscle strengthening activities should be done at least 3 days/week as part of the 60 or more minutes.
- Bone-strengthening activities should be done at least 3 days/week as part of the 60 or more minutes.

The physical activity guidelines for adults state:²³

- Adults should avoid inactivity. Some physical activity is better than none, and adults who participate in any amount of physical activity gain some health benefits.
- For substantial health benefits, adults should do at least 150 minutes (2 hours and 30 minutes) a week of moderate-intensity, or 75 minutes (1 hour and 15 minutes) a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate- and vigorous-intensity aerobic activity. Aerobic activity should be performed in episodes of at least 10 minutes, and preferably, it should be spread throughout the week.
- For additional and more extensive health benefits, adults should increase their aerobic physical activity to 300 minutes (5 hours) a week of moderate-intensity, or 150 minutes a week of vigorous-intensity aerobic physical activity, or an equivalent
- combination of moderate- and vigorous-intensity activity. Additional health benefits are gained by engaging in physical activity beyond this amount.
- Adults should also include muscle-strengthening activities that involve all major muscle groups on two or more days a week.

WHO Guidelines have easy to follow good practice statements for youth and adults with disabilities.

- For children with a disability ages 6-17 state:¹²
 - Doing some physical activity is better than doing none.
 - If children and adolescents living with disability are not meeting these recommendations, doing some physical activity will bring benefits to health.
 - Children and adolescents living with disability should start by doing small

Guidelines for the Care of People with Spina Bifida

amounts of physical activity and gradually increase the frequency, intensity and duration over time.

- There are no major risks for children and adolescents living with disability engaging in physical activity when it is appropriate to an individual's current activity level, health status and physical function; and the health benefits accrued outweigh the risks.
- Children and adolescents living with disability may need to consult a health-care professional or other physical activity and disability specialist to help determine the type and amount of activity appropriate for them.

For adults with a disability 18 years and above state:¹²

- Doing some physical activity is better than doing none.
- If adults living with disability are not meeting these recommendations, doing some physical activity will bring benefits to health.
- Adults living with disability should start by doing small amounts of physical activity, and gradually increase the frequency, intensity and duration over time.
- There are no major risks to adults living with disability engaging in physical activity when it is appropriate to the individual's current activity level, health status and physical function; and when the health benefits accrued outweigh the risks.
- Adults living with disability may need to consult a healthcare professional or other physical activity and disability specialist to help determine the type and amount of activity appropriate for them.

Please see Appendix for physical activity and disability resources.

Outcomes

Primary

1. Increase (or maintain) the minutes per day of daily physical activity at the different age levels as established by the National Physical Activity Guidelines.

Secondary

1. Increase knowledge and awareness of physical activity (i.e., benefits, safety, what/how to do it).

Tertiary

1. Improve health outcomes through physical activity participation across the lifespan (holistically, to maintain function, prevent secondary conditions, and for mental health and wellness).

0-11 months

Clinical Questions

Guidelines for the Care of People with Spina Bifida

1. Is there evidence that early motor skill intervention increases physical activity across the lifespan?
2. How early should doctors and therapists talk to parents/caregivers about physical activity for infants with Spina Bifida?
3. Are physical activity goals included in the Individual Family Service Plan (IFSP) for children beyond physical and occupational therapy?

Guidelines

1. Conduct infant motor development assessment to evaluate motor function in children with Spina Bifida to identify the most appropriate therapeutic intervention to enhance motor development outcomes.²⁴(Mobility Guidelines)
2. Provide guidance to parents and caregivers and include physical therapists in discussions about how to encourage movement and activity in their child.²⁵ (Health Promotion and Preventive Health Care Service Guidelines)
3. Inform parents and caregivers of their child's right to early intervention services that include adapted physical education/activity (Appendix: Early Intervention Services, Individualized Educational Plans (IEP) and 504 Plans). Encourage parents/caregivers to request that physical activity goals be added to the IFSP, if eligible for IFSP.²⁶

1-2 years 11 months

Clinical Questions

1. What strategies work to educate parents/caregivers about the importance of physical activity and ways to get their child involved?
2. How early should parent/caregiver education about physical activity start? What resources are available?
3. What is the most successful way to encourage parents/caregivers to invest time and money in physical activities for their child with Spina Bifida?

Guidelines

1. Discuss with parents and caregivers the benefits of involving their child with Spina Bifida in recreation, physical activity, and social programs and services, and provide information and/or resources about adapted and inclusive activities.²³ (Health Promotion and Preventive Health Care Services Guidelines)
2. Collaborate with parents/caregivers to identify physical activities they can do in everyday life to model the importance of physical activity as part of a healthy lifestyle. (clinical consensus)
3. Use motivational interviewing techniques with parents/caregivers to talk about physical activity goals for their child with Spina Bifida and work through barriers.²⁷
4. Inform parents/caregivers of the rights of their child to adapted physical education/activity and encourage parents/caregivers to advocate for physical activity goals to be added to their IFSP* or Section 504 plan (if they are eligible for an IFSP or Section 504 plan) (Appendix: Early Intervention Services, Individualized Educational Plans (IEP) and 504 Plans).²⁶

Guidelines for the Care of People with Spina Bifida

3-5 years 11 months

Clinical Questions

1. Given that some children with Spina Bifida will be limited in their physical function, what activities should they be exposed to, and when? Typically, children start playing baseball, soccer and other team sports at age 4 or 5. Should children with Spina Bifida start similar activities at the same age?
2. What training will coaches need to welcome children to meaningful play?
3. How does proper/improper mobility equipment affect the child's ability to participate in physical activity (e.g., a wheelchair that fits properly, proper seating and back setup, ankle foot orthotics, crutches, etc.)?
4. Who can parents and caregivers and doctors consult when they have questions related to physical activity and exercises that help maintain upper and lower body function (e.g., physical therapists (PTs), occupational therapists (OTs), recreation therapists, fitness staff, the National Center on Health, Physical Activity and Disability)?

Guidelines

1. Discuss with parents and caregivers the importance of involving the child in recreation, physical activity, limiting sedentary behaviors and engaging in social programs/services where they can be actively engaged with peers who have and those who don't have a disability.²⁸ Also, give parents/caregivers information about the life-long benefits of physical activity (e.g. active adults with Spina Bifida report more functional independence and a higher quality of life compared to those with Spina Bifida who are inactive).⁹ (Health Promotion and Preventive Health Care Services Guidelines)
2. Discuss strategies with parents/caregivers that balance parental involvement in their child's physical activities and the child's need for autonomy to increase independence.²⁹
3. Identify and provide additional support and information on precautions that children with shunts and ambulatory limitations should take when being physically active.³⁰
4. Use a team approach and include PTs/OTs to work with parents/caregivers to ensure children have proper fitting mobility equipment to maximize participation in physical activity. (clinical consensus)
5. Educate parents/caregivers of their child's right to adapted physical education/activity in preschool and encourage parents/caregivers to advocate for physical activity goals
6. to be added to their Individualized Education Plan (IEP) or Section 504 plan (if eligible for IEP or Section 504 plan).^{26,31}

6-12 years 11 months

Clinical Questions

1. What are some strategies to continue physical activity or physical education while in the hospital or after a long hospital stay (e.g., exercise band/tubing stretches or increasing knowledge of physical activity through reading/lecture- style learning if no physical activity is allowed)?
2. Is there evidence about the benefits/effects on the physical activity levels of children with Spina

Guidelines for the Care of People with Spina Bifida

- Bifida who advocate for themselves on their physical education goals in their IEP meetings?
3. Does the evidence support the least restrictive environment? Is this support related to health, socialization, something else, or a combination of factors?
 4. What training do schools need to include children with Spina Bifida in meaningful play throughout the day (at recess, physical education class, on-site after-school programs)?
 5. What are some ways that physical education teachers can be more inclusive of children with Spina Bifida? How do we start this process as early as possible?
 6. Does getting children engaged in sports at a young age improve the likelihood that they will remain engaged in activity throughout their lifespan?

Guidelines

1. Discuss the benefits of participating in physical activity, recreation, and sports with children with Spina Bifida.^{32,33} Discuss with parents/caregivers the importance of limiting sedentary behaviors.¹² Encourage parents to give their child choices about where they can be actively engaged with peers who have and those who don't have a disability.³⁴ (Health Promotion and Preventive Health Care Services Guidelines)
2. Recommend that parents/caregivers follow the National Physical Activity Guidelines for their child with Spina Bifida as closely as possible unless a health care provider advises that they are medically unsafe. (clinical consensus)
3. Engage in 60 minutes of physical activity or more each day.²³
4. Aerobic activity should make up most of the child's activity each day; vigorous intensity aerobic activity should be done at least 3 days/week.
5. Muscle strengthening activities should be done at least 3 days/week as part of the 60 or more minutes.
6. Bone strengthening activities should be done at least 3 days/week as part of the 60 or more minutes.
7. Use health care encounters and follow-up meetings to develop physical activity goals and monitor progress (e.g. minutes of physical activity per day). Use motivational interviewing techniques with the child and parents/caregivers to talk about physical activity goals and work through barriers.²⁷ Support parents/caregivers to develop an action plan with strategies to support their child's participation in physical activity in their community and school.³⁵ Prescribe, using a prescription pad, physical activity based on goals identified by the child.²³ (clinical consensus)
8. Perform pre-participation evaluations for children with Spina Bifida in collaboration with the child and family, pediatric specialists, therapists, coaches, and others to identify medical risks and modifications that can be made to ensure participation.³²
9. Identify strategies to minimize risk of illness and injury related to participation through activity adaptations and safety precautions.³² Identify and provide additional support and information for parents/caregivers on precautions to take when children with
10. shunts and ambulatory limitations are being active.³⁰ Discuss strategies that balance the parents'/caregivers' involvement with their child's need for independence when they participate in physical activity.²⁹
11. Use a team approach and include PTs/OTs to work with parents/caregivers to ensure their child has proper fitting mobility equipment to maximize physical activity participation. (clinical consensus)
12. Work with children with Spina Bifida and their family to address personal barriers such as

Guidelines for the Care of People with Spina Bifida

bowel/bladder care, medical events, assistive devices, as well as environmental factors that may affect participation.²¹

13. Advocate for and address barriers to participation of children with Spina Bifida in physical activity, recreation, and sports.³²
14. Inform parents/caregivers of their child's right to adapted physical education/activity and encourage parents/caregivers to advocate for physical activity goals to be added to their child's IEP or Section 504 plan (if eligible for IEP or Section 504 plan).^{26,31}
15. Advocate for the participation of children with Spina Bifida in both unified and adapted sports, recreation, and physical activity programs.³²
16. Provide families with a local/regional therapeutic recreation and adapted sport resource guide.³⁰

13-17 years 11 months

Clinical Questions

1. Do adapted physical education programs in schools adequately prepare (via a transition plan) children with Spina Bifida to lead physically active lifestyles?
2. What are the physical activity contraindications for children with Spina Bifida?
3. What are the types of physical activities used or recommended in the literature specific to children with Spina Bifida (resistance, cardio, incidental activity vs. planned physical activity/exercise)?
4. In what setting are children the most comfortable and likely to continue participation in a physical activity (e.g., in a group, at home, etc.)?
5. What are the doses of physical activity used or recommended in the literature for children with Spina Bifida? Are they effective for health changes?
6. Is physical activity included in the Individualized Transition Plan for children? (Transition Guidelines)

Guidelines

1. Discuss with children the benefits of participating in physical activity, recreation, and sports.^{32,35} Discuss with children and parents/caregivers the importance of limiting sedentary behaviors.²⁸ Encourage children and parents/caregivers to consider choices about where they can be actively engaged with peers who have and those who don't have a disability.³⁴ (Health Promotion and Preventive Health Care Services Guidelines)
2. Recommend that children and parents/caregivers follow the National Physical Activity Guidelines as closely as possible, unless a health care provider advises that they are medically unsafe. (clinical consensus)
 - Engage in 60 minutes of physical activity or more each day.²³
 - Aerobic activity should make up most of the child's activity each day; vigorous intensity aerobic activity should be done at least 3 days/week.
 - Muscle strengthening activities should be done at least 3 days/week as part of the 60 or more minutes.
 - Bone strengthening activities should be done at least 3 days/week as part of the 60

Guidelines for the Care of People with Spina Bifida

or more minutes.

3. Use health care encounters and follow-up meetings to develop physical activity goals and monitor progress (e.g. minutes of physical activity per day). Use motivational interviewing techniques with children and parents/caregivers to talk about physical activity goals and work through barriers.²⁷ Support children and parents/caregivers to develop an action plan with strategies to support the participation of children age 13- 17 years in physical activity in their community and school.³⁵ Prescribe, using a prescription pad, physical activity based on goals identified by children with Spina Bifida.²³ (clinical consensus)
4. Perform pre-participation evaluations for children with Spina Bifida in collaboration with parent/caregivers, pediatric specialists, therapists, coaches, and others to identify medical risks and modifications that can be made to ensure participation.³²
5. Identify strategies to minimize risks of illness and injury related to participation through activity adaptations and safety precautions.³² Identify and provide additional support and information for children age and parents/caregivers on precautions to take when children with shunts and ambulatory limitations are being active.³⁰ Discuss strategies that balance the parents'/caregivers' involvement with their child's need for independence when they participate in physical activity.²⁹
6. Use a team approach and include PTs/OTs to work with parents/caregivers to ensure their child has proper fitting mobility equipment to maximize physical activity participation. (clinical consensus)
7. Work with children with Spina Bifida and their family to address personal barriers such as bowel/bladder care, medical events, assistive devices, as well as environmental factors that may affect participation.²¹
8. Advocate for and address barriers to participation of children with Spina Bifida in physical activity, recreation, and sports.³²
9. Inform parents/caregivers of their child's rights to adapted physical education/activity and encourage children and parents/caregivers to advocate for physical activity goals to be added to their IEP or Section 504 plan (if eligible for IEP or Section 504 plan).^{26,31}
10. Assist students who are considering post-secondary education to assess supports for physical activity in the educational institutions they are considering and include these in their individual transition plans.
11. Advocate for the participation of children with Spina Bifida in both unified and adapted sports, recreation, and physical activity programs.³²
12. Provide families with a local/regional therapeutic recreation and adapted sport resource guide.³⁰
13. Discuss with children where they feel most comfortable being physically active and about their options (e.g., in the community, at school, with peers with/without disabilities).³⁴
14. Highlight that ages 13-17 years is a critical period to build physical activity into a daily routine that will preserve overall lifelong satisfaction and community participation

Guidelines for the Care of People with Spina Bifida

among persons with Spina Bifida.³³

18+ years

Clinical Questions

1. What are the physical activity contraindications for adults with Spina Bifida?
2. What are the types of physical activities used or recommended in the literature specific to adults with Spina Bifida (resistance, cardio, incidental activity vs. planned physical activity/exercise)?
3. What is the best setting for physical activity for adults (e.g., in a group, at home, etc.)? Where are they most comfortable and likely to continue participation?
4. What are the doses of physical activity used or recommended in the literature for adults with Spina Bifida? Are they effective for health changes?

Guidelines

1. Discuss the National Physical Activity Guidelines with adults with Spina Bifida.²³ (Health Promotion and Preventive Health Care Services Guidelines)
2. Follow the guidelines for adults with Spina Bifida as closely as possible, unless a health care provider advises that they are medically unsafe. (clinical consensus)
3. For substantial health benefits, it is recommended that adults should do at least 150 minutes (2 hours and 30 minutes) a week of moderate-intensity, or 75 minutes (1 hour and 15 minutes) a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate- and vigorous-intensity aerobic activity. It is recommended that aerobic activity should be performed in episodes of at least 10 minutes, and preferably, it should be spread throughout the week. (clinical consensus)
4. For additional and more extensive health benefits, it is recommended that adults should increase their physical activity to 300 minutes (5 hours) a week of moderate- intensity aerobic exercise, or 150 minutes (2 hours and 30 minutes) a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate- and vigorous-intensity activity. Additional health benefits are gained by engaging in physical activity beyond this amount. (clinical consensus)
5. It is recommended that adults should include muscle-strengthening activities that involve all major muscle groups on 2 or more days a week. (clinical consensus)
6. It is recommended that all adults should avoid inactivity. Some physical activity is better than none, and adults who participate in any amount of physical activity gain some health benefits. (clinical consensus)
7. Identify and provide additional support and information on physical activity precautions for adults with shunts and ambulatory limitations.³⁰
8. Use health care encounters and follow-up meetings to develop physical activity goals and monitor progress (e.g. minutes of physical activity per day). Employ motivational interviewing techniques to discuss and set physical activity goals and strategies to overcome barriers to achieving those goals.²⁷ Discuss the importance of physical activity and physical activity options

Guidelines for the Care of People with Spina Bifida

with adults with Spina Bifida.⁹

9. Prescribe, using a prescription pad, physical activity based off on goals discussed with adults with Spina Bifida.¹²
10. Assist students who are considering post-secondary education to assess supports for physical activity in the educational institutions they are considering.
11. Use a team approach and include PTs/OTs to work with the adult with Spina Bifida to make sure that their mobility equipment fits properly in order to maximize their participation in physical activities. (clinical consensus)
12. Emphasize that any movement is beneficial.¹²

Research Gaps

1. What are the strategies/solutions needed to intrinsically motivate individuals with Spina Bifida to be physically active at different ages across the lifespan (e.g., peer support, increased knowledge, self-efficacy, and other approaches)?
2. How can we best educate and train parents and individuals with Spina Bifida to be advocates for their inclusion in physical activity in their communities at large?
3. What physical activity resources are available for doctors nationwide? Locally? What resources need to be created?
4. Is there evidence that physical activity prevents secondary conditions? If so, how strong is this evidence?
5. What are the social/health benefits of participating in physical activity for individuals with Spina Bifida at different ages across the lifespan?
6. What are some strategies to continue physical activity or physical education while in the hospital or after a long hospital stay (e.g., exercise band/tubing stretches or increasing knowledge of physical activity through reading/lecture- style learning if no physical activity is allowed)?
7. Is there evidence that the early motor skill intervention increases physical activity across the lifespan?
8. What strategies work to educate parents/caregivers about the importance of physical activity and ways to get their child involved?
9. How early should parent/caregiver education about physical activity start? What resources are available?
10. What is the most successful way to encourage parents/caregivers to invest time and money in physical activities for their child with Spina Bifida?
11. Given that some children with Spina Bifida will be limited in their physical function, what activities should they be exposed to, and when? Typically, kids start playing baseball, soccer and other team sports at age 4-5 years. Should children with Spina Bifida start similar activities at the same age?
12. What training will coaches need to welcome children to meaningful play?
13. How does proper/improper mobility equipment affect the child's ability to participate in physical activity (e.g., a wheelchair that fits properly, proper seating and back set up, ankle foot orthotics, crutches, etc.)?
14. Who can parents/caregivers and doctors consult when they have questions related to physical activity/exercises that help maintain upper and lower body function (e.g., physical

Guidelines for the Care of People with Spina Bifida

- therapists (PTs), occupational therapists (OTs), recreation therapists, fitness staff, National Center on Health, Physical Activity and Disability)?
15. Is there evidence about the benefits/effects on the physical activity levels of children with Spina Bifida who advocate for themselves on their physical education goals in their IEP meetings?
 16. Does the evidence support the least restrictive environment? Is this support related to health, socialization, something else, or a combination of factors?
 17. What training do schools need to include children with Spina Bifida in meaningful play throughout the day (at recess, physical education class, on-site after-school programs)?
 18. What are some ways that physical education teachers can be more inclusive of children with Spina Bifida? How do we start this process as early as possible?
 19. Does getting children engaged in sports at a young age improve the likelihood that they will remain engaged in activity throughout their lifespan?
 20. What is the best setting for physical activity for adults (e.g., in a group, at home, etc.)? Where are they most comfortable and likely to continue participation?
 21. What are the doses of physical activity used or recommended in the literature for adults with Spina Bifida? Are they effective for health changes?
 22. What physical activity resources are available for doctors nationwide? Locally? What resources need to be created?
 23. Is there evidence that physical activity prevents secondary conditions? If so, how strong is this evidence?
 24. What are the social/health benefits of participating in physical activity for individuals with Spina Bifida at different ages across the lifespan?
 25. What are some strategies to continue physical activity or physical education while in the hospital or after a long hospital stay (e.g., exercise band/tubing stretches or increasing knowledge of physical activity through reading/lecture- style learning if no physical activity is allowed)?
 26. Is there evidence that the early motor skill intervention increases physical activity across the lifespan?
 27. What strategies work to educate parents/caregivers about the importance of physical activity and ways to get their child involved?
 28. How early should parent/caregiver education about physical activity start? What resources are available?
 29. What is the most successful way to encourage parents/caregivers to invest time and money in physical activities for their child with Spina Bifida?
 30. Given that some children with Spina Bifida will be limited in their physical function, what activities should they be exposed to, and when? Typically, kids start playing baseball, soccer and other team sports at age 4-5 years. Should children with Spina Bifida start similar activities at the same age?
 31. What training will coaches need to welcome children to meaningful play?
 32. How does proper/improper mobility equipment affect the child's ability to participate in physical activity (e.g., a wheelchair that fits properly, proper seating and back set up, ankle foot orthotics, crutches, etc.)?
 33. Who can parents/caregivers and doctors consult when they have questions related to physical activity/exercises that help maintain upper and lower body function (e.g., physical therapists (PTs), occupational therapists (OTs), recreation therapists, fitness staff, National

Guidelines for the Care of People with Spina Bifida

- Center on Health, Physical Activity and Disability)?
34. Is there evidence about the benefits/effects on the physical activity levels of children with Spina Bifida who advocate for themselves on their physical education goals in their IEP meetings?
 35. Does the evidence support the least restrictive environment? Is this support related to health, socialization, something else, or a combination of factors?
 36. What training do schools need to include children with Spina Bifida in meaningful play throughout the day (at recess, physical education class, on-site after-school programs)?
 37. What are some ways that physical education teachers can be more inclusive of children with Spina Bifida? How do we start this process as early as possible?
 38. Does getting children engaged in sports at a young age improve the likelihood that they will remain engaged in activity throughout their lifespan?
 39. What is the best setting for physical activity for adults (e.g., in a group, at home, etc.)? Where are they most comfortable and likely to continue participation?
 40. What are the doses of physical activity used or recommended in the literature for adults with Spina Bifida? Are they effective for health changes?

References

1. 2018 Physical Activity Guidelines Advisory Committee. *2018 Physical Activity Guidelines Advisory Committee Scientific Report*.; 2018.
2. Blair SN, Kohl HW, Paffenbarger RS, Clark DG, Cooper KH, Gibbons LW. Physical Fitness and All-Cause Mortality: A Prospective Study of Healthy Men and Women. *JAMA*. 1989;262(17):2395-2401. doi:10.1001/JAMA.1989.03430170057028
3. Cowan RE. Exercise Is Medicine Initiative: Physical Activity as a Vital Sign and Prescription in Adult Rehabilitation Practice. *Arch Phys Med Rehabil*. 2016;97(9):S232-S237. doi:10.1016/j.apmr.2016.01.040
4. Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines Advisory Committee Report, 2008. To the Secretary of Health and Human Services. Part A: Executive Summary*.; 2008.
5. Zenko Z, Willis EA, White DA. Proportion of Adults Meeting the 2018 Physical Activity Guidelines for Americans According to Accelerometers. *Front Public Health*. 2019;7:135-135. doi:10.3389/fpubh.2019.00135
6. van den Berg-Emons HJG, Bussmann JBJ, Meyerink HJ, Roebroek ME, Stam HJ. Body fat, fitness and level of everyday physical activity in adolescents and young adults with meningomyelocele. *J Rehabil Med*. 2003;35(6):271-275. doi:10.1080/16501970310012400
7. Schoenmakers MAGC, de Groot JF, Gorter JW, Hillaert JLM, Helden PJM, Takken T. Muscle strength, aerobic capacity and physical activity in independent ambulating children with lumbosacral spina bifida. *Disabil Rehabil*. 2009;31(4):259-266. doi:10.1080/09638280801923235
8. van den Berg-Emons HJ, Bussmann JB, Brobbel AS, Roebroek ME, van Meeteren J, Stam HJ. Everyday physical activity in adolescents and young adults with meningomyelocele as

Guidelines for the Care of People with Spina Bifida

- measured with a novel activity monitor. *J Pediatr*. 2001;139(6):880-886. doi:10.1067/mpd.2001.119991
9. Rimmer JH. Exercise and physical activity in persons aging with a physical disability. *Phys Med Rehabil Clin N Am*. 2005;16(1):41-56. doi:10.1016/j.pmr.2004.06.013
 10. Buffart LM, Roebroek ME, Rol M, Stam HJ, van den Berg-Emons RJG. Triad of physical activity, aerobic fitness and obesity in adolescents and young adults with myelomeningocele. *Acta Derm Venereol*. 2008;40(1):70-75. doi:10.2340/16501977-0135
 11. Buffart LM, Van Den Berg-Emons RJG, Meeteren J Van, Stam HJ, Roebroek ME. Lifestyle, participation, and health-related quality of life in adolescents and young adults with myelomeningocele. *Dev Med Child Neurol*. 2009;51(11):886-894. doi:10.1111/j.1469-8749.2009.03293.x
 12. WHO Guidelines on Physical Activity and Sedentary Behaviour.; 2020.
 13. Ding D, Mutrie N, Bauman A, Pratt M, Hallal PRC, Powell KE. Physical activity guidelines 2020: comprehensive and inclusive recommendations to activate populations. *The Lancet (British edition)*. 2020;396(10265):1780-1782. doi:10.1016/S0140-6736(20)32229-7
 14. Carty C, van der Ploeg HP, Biddle SJH, et al. The First Global Physical Activity and Sedentary Behavior Guidelines for People Living With Disability. *J Phys Act Health*. 2021;18(1):86-93. doi:10.1123/jpah.2020-0629
 15. Martin Ginis KA, Latimer-Cheung AE, West CR. Commentary on “The First Global Physical Activity and Sedentary Behavior Guidelines for People Living With Disability.” *J Phys Act Health*. 2021;18(4):348-349. doi:10.1123/jpah.2020-0871
 16. Carty C, van der Ploeg HP, Biddle SJH, et al. Response to “Commentary on: The First Global Physical Activity and Sedentary Behavior Guidelines for People Living With Disability.” *J Phys Act Health*. 2021;18(4):350-351. doi:10.1123/jpah.2021-0137
 17. Ginis KAM, Hicks AL, Latimer AE, et al. The development of evidence-informed physical activity guidelines for adults with spinal cord injury. *Spinal Cord*. 2011;49(11):1088-1096. doi:10.1038/sc.2011.63
 18. Oliveira A, Jácome C, Marques A. Physical fitness and exercise training on individuals with Spina Bifida: A systematic review. *Res Dev Disabil*. 2014;35(5):1119-1136. doi:10.1016/j.ridd.2014.02.002
 19. Monti R, Mariani F, Masticci R, et al. Spina bifida and cardiorespiratory profile: the impact of leisure sport activities on physical fitness. *Child's nervous system*. 2024;40(1):205-211. doi:10.1007/s00381-023-06152-3
 20. Lai B, Davis D, Narasaki-Jara M, et al. Feasibility of a Commercially Available Virtual Reality System to Achieve Exercise Guidelines in Youth With Spina Bifida: Mixed Methods Case Study. *JMIR Serious Games*. 2020;8(3):e20667-e20667. doi:10.2196/20667
 21. Bloemen MAT, Verschuren O, van Mechelen C, et al. Personal and environmental factors to consider when aiming to improve participation in physical activity in children with Spina Bifida: A qualitative study. *BMC Neurol*. 2015;15(1). doi:10.1186/s12883-015-0265-9
 22. Bloemen M, Takken T, Groot J, et al. Determinants of physical activity in young wheelchair-user with spina bifida. *J Rehabil Med*. 2020;52(10):jrm00115-jrm00115. doi:10.2340/16501977-2724
 23. U.S. Department of Health and Human Services. *Physical Activity Guidelines for Americans 2nd Edition*.; 2018.
 24. Darrah J, Piper M, Watt M. Assessment of gross motor skills of at-risk infants: predictive validity of the Alberta Infant Motor Scale. *Dev Med Child Neurol*. 1998;40(7):485-491.

Guidelines for the Care of People with Spina Bifida

- doi:10.1111/j.1469-8749.1998.tb15399.x
25. Adolph KE, Franchak JM. The development of motor behavior. *Wiley Interdiscip Rev Cogn Sci.* 2017;8(1-2):n/a. doi:10.1002/wcs.1430
 26. Dummer GM, Connor-Kuntz FJ, Goodway JD. A Physical Education Curriculum for All Preschool Students. *Teach Except Child.* 1995;27(3):28-34. doi:10.1177/004005999502700307
 27. Resnicow K, Davis R, Rollnick S. Motivational Interviewing for Pediatric Obesity: Conceptual Issues and Evidence Review. *J Am Diet Assoc.* 2006;106(12):2024-2033. doi:10.1016/j.jada.2006.09.015
 28. Law M, King G, King S, et al. Patterns of participation in recreational and leisure activities among children with complex physical disabilities. *Dev Med Child Neurol.* 2006;48(5):337-342. doi:10.1017/S0012162206000740
 29. Antle BJ, Mills W, Steele C, Kalnins I, Rossen B. An exploratory study of parents' approaches to health promotion in families of adolescents with physical disabilities. *Child Care Health Dev.* 2008;34(2):185-193. doi:10.1111/j.1365-2214.2007.00782.x
 30. Flanagan A, Gorzkowski M, Altiock H, Hassani S, Ahn KW. Activity Level, Functional Health, and Quality of Life of Children with Myelomeningocele as Perceived by Parents. *Clin Orthop Relat Res.* 2011;469(5):1230-1235. doi:10.1007/s11999-010-1651-7
 31. Murata NM, Maeda JK. Structured Play for Preschoolers with Developmental Delays. *Early Child Educ J.* 2002;29(4):237. doi:10.1023/A:1015181607622
 32. Murphy NA, Carbone PS, and Council on Children With Disabilities. Promoting the Participation of Children With Disabilities in Sports, Recreation, and Physical Activities. *Pediatrics.* 2008;121(5):1057-1061. doi:10.1542/peds.2008-0566
 33. Dykens EM, Rosner BA, Butterbaugh G. Exercise and Sports in Children and Adolescents with Developmental Disabilities: Positive Physical and Psychosocial Effects. *Child Adolesc Psychiatr Clin N Am.* 1998;7(4):757-771. doi:10.1016/S1056-4993(18)30210-4
 34. Knibbe TJ, Biddiss E, Gladstone B, McPherson AC. Characterizing socially supportive environments relating to physical activity participation for young people with physical disabilities. *Dev Neurorehabil.* 2017;20(5):294-300. doi:10.1080/17518423.2016.1211190
 35. Rimmer JA, Rowland JL. Physical activity for youth with disabilities : A critical need in an underserved population. *Phys Ther.* 2008;11(June):141-148. doi:10.1080/17518420701688649

Appendix

- National Physical Activity Guidelines for Americans, Office of Disease Prevention and Health Promotion, Office of the Assistant Secretary for Health, Office of the Secretary, U.S. Department of Health and Human Services
- The Centers for Disease Control and Prevention's disability & physical activity resources for doctors and other health professionals, CDC Disability and Health Branch
- The National Center on Health, Physical Activity, and Disability (NCHPAD):
- United States Olympic Committee Para Sport
- World Health Organization Physical Activity